



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Genesis of the Zinc Ores of Edwards District, St. Lawrence County, N.Y. By C. H. SMYTH, JR. New York State Museum Bulletin, No. 201. Albany, N.Y., 1917. Pp. 39, pls. 12.

This report presents the results of laboratory studies of ore from the zinc deposits of the Edwards district, which is about 15 miles south of the Canton area described in Bulletin 185, reviewed above, and the geology of the two areas is very similar. The ore minerals are sphalerite, pyrite, and galena, and occur as fillings of narrow cracks or as replacements along shear zones in the Grenville crystalline limestone. The deposits are of the high-temperature type, and the sequence of mineral deposition is as follows: (1) diopside, tremolite, (2) pyrite, (3) sphalerite, (4) galena, (5) talc, (6) serpentine. This sequence of mineral deposition indicates changing conditions from intense-contact, metamorphic conditions to those normal for the depths involved and without outside agencies. At all stages during this transition from intense to moderate conditions calcite was subject to repeated solution and recrystallization.

The granite magmas which intrude the Grenville sediments are considered the source of the ore minerals and during the cooling of the magma, gases and solutions were given off which carried the sulphur and metals into the country rocks where they were precipitated. The wall rock of the sphalerite is always crystalline limestone while the typical pyrite ore is always in schist or gneiss. Also the pyrite ore is always rich in graphite while the zinc ore contains but little graphite. The typical pyrite ore contains but little sphalerite but the zinc ore contains considerable pyrite.

These deposits are compared with zinc deposits of other regions and especially with the contact zinc deposits of the Christiania district described by Goldschmidt. Excellent photomicrographs show clearly the various mineral relations described in detail in the text. This is a valuable contribution to the subject of contact metamorphic ore deposits.

J. F. W.

New Edition of Coal, Oil, Gas, Limestone, and Iron Ore Map. West Virginia Geological Survey.

Thoroughly revised, showing oil and gas pools, many anticlinal lines not heretofore shown, and also booklet giving the names and post-office addresses of all the principal coal-mining operators in West Virginia up to July 1, 1921. Scale, 8 miles to the inch. Price, folded in strong envelope and delivered by mail, \$1.00. Remittances to West Virginia Geological Survey, Box 848, Morgantown, West Virginia.